

VFC - Toe fracture (not big toe)

Toe fracture (not big toe)

You have a fracture of your toe (phalanx fracture). This injury is treated with a tape, called a buddy tape.



Summary

- You have a fracture of your toe.
- This type of injury almost always heals well without any loss of function.
- A routine follow-up is not necessary for this type of injury.
- You can remove the buddy tape after 1-2 weeks as soon as the pain allows. Wear a sturdy shoe for 2 weeks.
- To prevent the toe from getting stiff, it is important that you start moving the toe again as soon as possible.
- You can find instructions on how to reapply the buddy tape in the app.
- If you have any questions or problems during the recovery, do not hesitate to contact us on workdays from 09:00 – 11:00 AM – 2:00 – 4:00 PM via the Fracture Hot Line.

Treatment

A fracture of the toe is a common injury. The fracture is treated with a buddy tape. A buddy tape attaches the injured toe to the toe next to it, immobilizing the injured toe.

0-1 week
<ul style="list-style-type: none"> • Buddy tape and a sturdy shoe.
1-2 weeks
<ul style="list-style-type: none"> • Buddy tape, remove it as soon as your pain allows this. • Wear a sturdy shoe.
After 3 weeks
<ul style="list-style-type: none"> • Start sports.

Instructions

Follow these instructions for a good recovery.

- Wear the buddy tape for 1-2 weeks. If necessary, you can replace the buddy tape. You can buy new tape at the pharmacy or use the tape that you received at the Emergency Department. In the app you can find instructions on how to reapply the buddy tape.
- You can stand on your foot if your pain allows you to do so. If necessary, you can rent crutches at the hospital.
- Elevate your foot when you are resting to reduce the swelling and pain. This can be done by putting your foot on a chair.
- To prevent the toe from getting stiff, it is important that you start moving the toe again as soon as possible.
- We advise you not to practise sports the first three weeks. After three weeks you can start sports again, if the pain allows you to do this.
- Are you in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait with the use of paracetamol until the pain is too much.

Recovery

- This type of injury almost always heals well with time and use – no specific treatment is required and routine follow-up is not necessary.
- Does the pain increase, or does it not improve? Please contact the Fracture Hot Line. If a check-up seems necessary, we will schedule an appointment.

Reapply buddy tape

If you want to reapply the buddy tape, you can watch the instructional video below. You can buy new materials at the pharmacy or reapply the tape from the Emergency Department.

https://www.youtube.com/watch?v=_vn41BOIZSs

In the video a special shoe is used, usually you do not need this special shoe and you can use your own (sturdy) shoe.

Exercises

It is important to move your toe as much as possible. No specific exercises are necessary.

Any questions?

Do you have any questions after reading this information? Ask your question via Laurentius hospital Fracture Hot Line.

Virtual Fracture Care App

Find more information about your injury in our app. The app also includes exercises that promote your recovery. You can find the app in the App Store (iPhone) or Google Store (Android).



You can also scan the QR code. You will then automatically be redirected to the app.

Contact

Fracture Hot Line

T 0475-38 22 11
(business days from 09:00 – 11:00 AM – 2:00 – 4:00 PM)

Source brochure: Proefschrift Virtual Fracture Care: Direct ontslag vanaf de SEH voor patiënten met simpele stabiele letsels

Laurentius Ziekenhuis
Mgr. Driessenstraat 6
(hoofdingang bereikbaar via de Oranjelaan)
6043 CV Roermond
www.lzr.nl

Patiëntencommunicatie Laurentius Ziekenhuis
VFC13 - Spoedeisende Hulp - oktober 2024